

Break O'Day National Park

Natural values of a proposed new National Park in North East Tasmania



August 2011

Executive Summary

The Break O'Day National Park seeks to recognise and protect the outstanding biodiversity values and landscapes of the St Marys region in northeast Tasmania. The National Park would connect several smaller existing reserves resulting in a single reserve which covers a wide range of environmental variation from near-coastal to elevated peaks.

Conservation values of the proposed Break O'Day National Park include:

- A diverse variety of different wet and dry forest types including old growth forests;
- Key habitat for one of Tasmania's most threatened invertebrates, the blind velvet worm (*Tasmanipatus anophthalmus*), and the related rare giant velvet worm (*Tasmanipatus barretti*);
- Nesting habitat for threatened bird species including swift parrot (*Lathamus discolor*), grey goshawk (*Accipiter novae-hollandiae*) and wedge-tailed eagle (*Aquila audax fleayi*);
- Brookers gum (*Eucalyptus brookeriana*) forest – an endangered vegetation type which is very rare in eastern Tasmania;
- Threatened plant species such as the Guinea flower (*Hibbertia calycina*) and Roundleaf mintbush (*Prostanthera rotundifolia*) – both shrubs listed as vulnerable to extinction;
- High biogeographic significance based on the distribution of flora and fauna species including several National Estate areas recognising the unique biogeography of the area;
- An exceptional diversity of lichens, including several species rare or not known from elsewhere in Tasmania;
- Numerous significant landforms which are recognised geoconservation sites including Huntsmans Creek Waterfall, Upper Durham Creek Karst, St Marys Porphyrite and Catos Creek Dyke.

The unique and exceptional values of this area warrant a higher degree of protection than the existing small Forest Reserves and State Reserves. The biological significance of the area and the landscape features (scenery, water catchments, geoconservation values) are sufficient to justify National Park status. Management of a single integrated reserve would help maintain the integrity and viability of these important values.

Report prepared by Nick Fitzgerald, BSc(Hons)

313 Nelson Rd, Mount Nelson, Tasmania

TasBotanic@gmail.com

August 2011

Cover photo: dry eucalypt forest recovering from fire on the steep slopes of North Sister.

Introduction

This report outlines the outstanding natural values of the proposed Break O'Day National Park (see map p. 20). Other values which contribute to the conservation status of this area including cultural heritage, tourism and landscape aesthetics are not considered here.

The hills around St Marys rise steeply from the East Coast providing a connection between the coastal environment, the inland grasslands of the Fingal Valley, the dolerite highlands to the south and the granite hills to the north. This area is a mixing zone of different landscape types, geologies, climatic zones, species and ecological communities.

Existing reserves in the area are small to medium sized, managed by different agencies and separated by other land tenures. Substantial areas with high conservation values have only informal reserve status.

Connecting the Avenue River Forest Reserve north of St Marys to the Douglas Apsley National Park to the south will result in a substantial corridor of protected land in the coastal hinterland of Tasmania's East Coast. There are no other options for preserving native vegetation connections between the Eastern Tiers and North East Highlands due to land clearing in the Fingal Valley. It is also crucial to connect the coastal environment with the hinterland which rises steeply inland and this will require initiatives involving private landowners in the coastal strip.

This connectivity is important because it allows the continuity of native habitats along environmental gradients (e.g. in altitude and distance from the coast) and provides a corridor for animals and plants to move through the landscape over short and long time scales (McQuillan *et al.* 2009).

Management as a National Park will also protect upper catchment areas which are important for ensuring water quantity and quality for lower catchments including coastal wetlands and estuaries such as Hendersons Lagoon as well as providing water for St Marys and other human settlements in the area.

Land Tenure

The proposed National Park covers around 11 664 hectares of public land, comprising seven existing reserves under the *Nature Conservation Act* and the *Forestry Act* (Table 1), plus areas of State forest and Crown Land (see map p. 21). Existing formal reserves account for 5703 hectares of the total area with the remaining 5 961 ha comprising State forest and Crown Land.

Table 1. Existing reserves within the proposed Break O'Day National Park.

Existing reserve	Size (hectares)	Gazettal date	Management authority
Nicholas Range Forest Reserve	810	18/12/1998	Forestry Tasmania
St Marys Pass State Reserve	367	30/03/1950	Parks & Wildlife Service
St Patricks Head State Reserve	1339	5/09/1963	Parks & Wildlife Service
German Town Forest Reserve	935	18/12/1998	Forestry Tasmania
Huntsmans Cap Forest Reserve	216	18/12/1998	Forestry Tasmania
Little Beach State Reserve	945	29/04/1995	Parks & Wildlife Service
Lower Marsh Creek Forest Reserve	1091	5/01/2005	Forestry Tasmania



Figure 1. Aerial photograph mosaic showing land tenure in the St Marys region, north east Tasmania.

The proposed National Park is contiguous with several areas of private land managed for nature conservation through legal instruments such as conservation covenants or Private Nature Reserves. A National Park would complement these private land conservation initiatives and together contribute to a wholistic landscape-scale plan for conservation and complementary land uses as envisioned by the St Marys Protected Landscape concept (NETLT 2009). There are at least ten permanent private land protected areas totalling more than 340 hectares in the St Marys region, which is an exceptional commitment to conservation from private landowners.

Boundary Justification

The southern part of the proposed National Park comprises the Lower Marsh Creek FR, Little Beach, St Patricks Head and St Marys State Reserves plus all of the adjoining State Forest. Effectively it includes all of the public land in this area, since this is needed to achieve a well connected and buffered reserve at the landscape scale. To the east of the German Town Forest Reserve an area of State Forest has been included so as to increase the representation of lowland dry eucalypt forest including oldgrowth in the proposed National Park and to improve connectivity toward the coast and the north. State Forest around the Nicholas Range FR and Huntsmans Cap FR is included to connect existing reserves, including German Town FR to the east and Avenue River FR to the north. These areas of State Forest include significant patches of

oldgrowth forest. The northern boundary of the proposed National Park is largely determined by the presence of pine plantations and extensive areas of silvicultural regrowth eucalypt forest. The lower slopes of the Nicholas Range include oldgrowth forest amongst silvicultural regrowth and consequently the whole public forest estate in these areas has been included to achieve integrity for the National Park. A Crown Land parcel at Cheeseberry Hill is included because it contains an important stand of *Eucalyptus brookeriana* wet forest.

By increasing the size of the protected area and consequently the variety of landscape features included and bringing the whole area under a single management plan the opportunities for biodiversity to adapt to climate change are increased.

Description of the proposed Break O’Day National Park

Climate

The region has a mild maritime climate at the coast grading rapidly into a more continental climate west of the drainage divide where the higher elevation results in colder minimum temperatures.




Figure 2. Mean monthly rainfall at St Marys (Lat: 41.58°S Lon: 148.18°E, Elevation: 258 m, records: 1905-2010) and Falmouth (Lat: 41.51°S Lon: 148.26° E, Elevation: 6 m, records: 1967-2010). Source: BOM.

Rainfall data from the coast at Falmouth show relatively constant rainfall throughout the year with an annual average of 666 mm while St Marys at 258 metres elevation and 9 km from the coast has a winter rainfall peak and an annual average of 1015 mm (Figure 2). Easterly weather patterns can bring intense rainfall to this region resulting in some of the heaviest rainfall events in Tasmania, particularly at higher elevations. For example, in the 12 years from 1999 to 2010, Gray near Mount Elephant has recorded 24 hour rainfall greater than 100 mm on 18 days and monthly totals of over 500 mm on two occasions, including in a single 4 day rainfall event. Cloud frequently gathers on the higher peaks such as St Patricks Head and consequently fog drip precipitation may add to rainfall and increase humidity at these elevations.

The nearest temperature records to the proposed NP are at Scamander (on the coast 3.5 km to the NW) and Fingal (11-25 km inland from the proposed NP). Monthly and annual maximum temperatures are very similar between the Scamander and Fingal (18°C and 17.8°C respectively, Figure 3). However Fingal has significantly lower minimum temperatures with an annual average of 5.3°C compared to 8.8°C at Scamander. Frosts occur frequently in winter around St Marys.




Figure 3. Mean monthly temperature at sites near the proposed National Park – Scamander (Latitude: 41.46°S Longitude: 148.26°E, Elevation : 3 m, records: 1974-2010) and Fingal (Latitude: 41.64°S Longitude: 147.97°E, Elevation : 237 m, 27 km from coast, records: 1882-2010).

Geology

Geologically the proposed National Park covers three major rock types (see map p. 22). The western part of the area is largely weathered Jurassic dolerite, with exposed dolerite on the peaks of the Nicholas Range, Mt Durham, Huntsmans Cap and St Patricks Head. Ordovician sedimentary rock (“Mathinna beds”) occur at lower elevations at the northern edges of the area. Most of the eastern portion of the area is composed of Devonian porphyry with an outcrop of dolerite at St Patricks Head. Small areas of Permian sedimentary rocks appear in several widespread locations, including the lower slopes of Mt Elephant. Limestone occurs in some of these areas, with associated karst features occurring at the Nicholas Range and Mt Elephant. Karst is rare in the northeast and east coast. The Upper Durham Creek Karst is of regional geoconservation significance and is vulnerable to changes in catchment land use.

A variety of soils occur in the area, reflecting the different bedrock types and the landscape processes and geological history.

Landscape and drainage

The varied and complex landscape of the proposed National Park creates three major drainage patterns. The southern slopes of the Nicholas Range and the western slopes of Mount Elephant drain inland, with creeks feeding the Break O’Day River which flows west to join the South Esk River. Consequently rain falling only 6 kilometres inland from the east coast can end up flowing over 200 kilometres west and north to eventually reach the Tamar Estuary and Bass Strait. The northern part of the National Park features north-flowing creeks which enter the Avenue and

Scamander rivers. The eastern slopes of the protected area comprise a series of small catchments with deeply incised creeks flowing east directly to the coast.

Vegetation

Eucalypt forest is the major vegetation type in the area (see map p. 23). The St Marys region features a high diversity of eucalypt species compared to elsewhere in north east Tasmania and several distinct forest types are recognised. Vegetation in this area is determined largely by moisture availability and soil fertility. The variation in altitude also influences vegetation patterns.

Dry open forests dominated by Ironbark (*Eucalyptus sieberi*) occur on dry sites on nutrient-poor soils throughout much of the northern and eastern parts of BODNP. Other dry forest types are characterised by either black peppermint (*E. amygdalina*) or stringybark (*E. obliqua*) occur in sizable patches. Wet stringybark forests grow on more sheltered sites such as gullies and south-facing slopes. At higher altitudes on the Nicholas Range gum-topped stringybark (*E. delegatensis*) replaces stringybark as the canopy species. There are only small areas of non-forest vegetation such as wet heath and tea-tree scrub.

Bioregions

The proposed national park occurs at the boundary of three of Tasmania's nine bioregions, representing the exceptional convergence of different environments and species in this area. Most of the proposed national park is within the easternmost part of the Ben Lomond bioregion, while the lower elevation near-coastal hills are in the southernmost extent of the Flinders bioregion. The northern boundary of the South East bioregion is at the southern edge of the proposed reserve.

Natural Values

Threatened Flora

Twenty-one threatened flora species are presently known from the proposed National Park (Table 2, map p. 27), however some areas have not been adequately surveyed and as such there are likely to be additional threatened species present, such as those known from the surrounding area (Table 3, map p. 27).

Table 2. Threatened flora recorded from Proposed National Park.

Name	Common Name	Tas status	National status	Distribution in Proposed National Park
<i>Asplenium trichomanes</i> subsp. <i>trichomanes</i>	dolerite spleenwort	v		Recorded from Mount Durham and Huntsman Cap in the proposed NP. Occupies dolerite outcrops.
<i>Blechnum cartilagineum</i>	gristle fern	v		Occurs in riparian situations along Little Beach Creek and Tin Creek in the proposed NP and on neighbouring private land. These catchments are key populations for the species in Tasmania.
<i>Cyathea cunninghamii</i>	slender treefern	e		Largest populations in Tasmania occur in sheltered gullies in Little Beach Creek and Lower Marsh Creek catchments in the proposed NP.
<i>Cyathea Xmarcescens</i>	skirted treefern	e		Largest populations in Tasmania occur in sheltered gullies in Little Beach Creek and Lower Marsh Creek catchments in the proposed NP.
<i>Desmodium gunnii</i>	southern ticktrefoil	v		Numerous records within and near the proposed NP, including in Little Beach SR and German Town FR. Occurs in dry forest, particularly near the East Coast.
<i>Deyeuxia densa</i>	heath bentgrass	r		Recorded from Huntsmans Cap, Nicholas Range and Lower Marsh Creek FR. Occurs in a variety of open habitats.
<i>Euphrasia collina</i> subsp. <i>deflexifolia</i>	eastern eyebright	r		Numerous records from State Forest around South Sister. Occurs in open woodland, dry forest and heath near the East Coast.
<i>Glycine microphylla</i>	small-leaf glycine	v		Recorded from German Town FR and Little Beach SR, also from several locations near the coast to the east of the proposed NP. Occurs in dry forest.
<i>Hibbertia calycina</i>	lesser guineaflower	v		Occurs near McIntyre Creek in the north of the proposed NP. This is one of the key sites for this species which is restricted to dry north-facing slopes in <i>Eucalyptus sieberi</i> forest near Scamander and St Helens.

<i>Hierochloa rariflora</i>	cane holygrass	r		Recorded from Lower Marsh Creek FR and German Town FR. Likely to be more widespread in dry and damp forest below around 400 m elevation in the proposed NP. Also known from this habitat in sites near the proposed NP.
<i>Hovea corrickiae</i>	glossy purplepea	r		Recorded from German Town FR, which is the largest of 33 known populations of this species. Also occurs at Mt Elephant and Lower Marsh Creek FR. Confined to riparian sites and damp slopes in the northeast.
<i>Hovea tasmanica</i> (previously <i>H. longifolia</i>)	rockfield purplepea	r		Known from State Forest on the northern slopes of the Nicholas Range. Occurs on dry rocky or riparian sites in open forest.
<i>Pellaea calidirupium</i>	hotrock fern	r		Recorded from dolerite outcrops at Mt Durham and Huntsmans Cap. Potentially present on other dolerite outcrops in the proposed NP.
<i>Phebalium daviesii</i>	davies waxflower	e	CR	Naturally confined to the George River near St Helens. An ex-situ population has been established in the St Patricks Head SR. Extremely rare riparian species.
<i>Plantago debilis</i>	shade plantain	r		Numerous records from various locations. Extensive potential habitat in both wet and dry forest.
<i>Pomaderris intermedia</i>	lemon dogwood	r		Recorded from Elephant Pass, which is considered a key site for this species. Potentially more widespread in wet forest in the area.
<i>Prostanthera rotundifolia</i>	roundleaf mintbush	v		Known from German Town FR and St Marys Pass SR. Common along the Scamander River north of the proposed NP. Occurs on dry hillsides and riverbanks. Estimated only 12 populations in Tasmania.
<i>Pterostylis ziegeleri</i>	grassland greenhood	v	VU	Recorded from State Forest near sea level close to Lagoons Beach. Coastal species unlikely to occur elsewhere in the proposed NP.
<i>Teucrium corymbosum</i>	forest germander	r		Known from St Marys Pass, St Patricks Head and Lower Marsh Creek FR. Occurs in a variety of habitats.
<i>Thismia rodwayi</i>	fairy lanterns	r		Occurs on southern slopes of South Sister on State Forest and private land. May occur elsewhere in suitable habitat: under leaf litter of wet forest understorey species on dolerite soils.
<i>Veronica plebeia</i>	trailing speedwell	r		Recorded from St Patricks Head. Potentially more widespread in wet forest.

Threatened Species Protection Act 1995 (Tasmania): r = rare, v = vulnerable, e = endangered, x = extinct

Environmental Protection and Biodiversity Conservation Act 1999 (Commonwealth): VU = vulnerable, EN = endangered, CR = critically endangered

Table 3. Threatened flora recorded from within 5 km and possibly present within Proposed National Park.

Name	Common Name	Tas status	National status	Location of known records and potential habitat in proposed National Park
<i>Acacia siculiformis</i>	dagger wattle	r		Suitable habitat is present in dry forest. Known from a site west of the proposed NP.
<i>Acacia ulicifolia</i>	juniper wattle	r		Suitable habitat is present in dry forest near the coast. Several records from north of the proposed NP near Scamander.
<i>Asperula subsimplex</i>	water woodruff	r		Potential habitat occurs in wetlands and riverbanks. Recorded from north of the proposed NP near Scamander.
<i>Austrostipa nodosa</i>	knotty speargrass	r		Suitable habitat is present in dry forest. Recorded from varied habitats and elevations north and south of the proposed NP at the Scamander River and Fingal Tier.
<i>Caladenia caudata</i>	tailed spider-orchid	v	VU	Recorded from low elevation sites near Lagoons Beach just east of the proposed NP. Limited potential habitat in proposed NP, prefers sandy and loamy soils in heathland or open woodland.
<i>Caladenia congesta</i>	blacktongue finger-orchid	e		Recorded from private land west of Mt Elephant just outside of the proposed NP. Potential habitat in dry open forest, particularly sandy or loamy soils.
<i>Conospermum hookeri</i>	tasmanian smokebush	v	VU	Recorded from coastal location near Scamander north of the proposed NP. Limited potential habitat occurs in low elevation heathy woodland in the proposed NP.
<i>Cynoglossum australe</i>	coast houndstongue	r		Generally a coastal species but does occur occasionally inland. Potential habitat in dry forests. Known from coastal sites at Scamander and within 250 m of the southern end of the proposed NP in Lagoons Beach Coastal Reserve.
<i>Hibbertia virgata</i>	twiggy guineaflower	r		Occurs near the coast south of Scamander. Prefers sandy coastal heath and near-coastal woodlands. Unlikely to occur in the proposed NP.
<i>Juncus amabilis</i>	gentle rush	r		Recorded from Tasman Highway under 1km east of proposed NP. Often found in roadside seepage areas. Potential habitat occurs in the proposed NP.
<i>Lepidium hyssopifolium</i>	soft peppergrass	e	EN	Known from roadside of Falmouth Road around 2 km east of proposed NP. Occurs in grasslands, grassy woodlands and on roadsides. Limited potential habitat in proposed NP.

<i>Lepidium pseudotasmanicum</i>	shade peppergrass	r		Known from roadside of Falmouth Road around 2 km east of proposed NP. Occurs in grasslands, grassy woodlands and on roadsides. Limited potential habitat in proposed NP.
<i>Lepidosperma tortuosum</i>	twisting rapier sedge	r		Several records from Tasman Highway south of St Marys around 1.5 km west of proposed NP at Mount Elephant. Some potential habitat in proposed NP in open woodland.
<i>Olearia hookeri</i>	crimsontip daisybush	r		Known from dry forest 3 km south of proposed NP. Potential habitat occurs on dry rocky sites in open forest.
<i>Pimelea curviflora</i>	curved rice-flower	r		Recorded from hills southwest of the proposed NP. Potential habitat in wet forest.
<i>Pterostylis atriola</i>	snug greenhood	r	EN	Known from Thompsons Marshes in hills 3 km south of proposed NP. Potential habitat in shrubby open eucalypt forest.
<i>Scutellaria humilis</i>	dwarf skullcap	r		Recorded from near Four Mile Creek east of proposed NP. Potential habitat in wet shady areas.
<i>Tricoryne elatior</i>	yellow rushlily	v		Recorded from near Four Mile Creek east of the proposed NP. Limited suitable habitat in open woodland.
<i>Xerochrysum palustre</i>	swamp everlasting		VU	Known from Thompsons Marshes in hills 3 km south of proposed NP. Limited potential habitat in swampy or riparian vegetation.

Threatened Species Protection Act 1995 (Tasmania): r = rare, v = vulnerable, e = endangered, x = extinct
Environmental Protection and Biodiversity Conservation Act 1999 (Commonwealth): VU = vulnerable, EN = endangered, CR = critically endangered

Lower Marsh Creek is a hotspot for rare and threatened ferns (Garrett 1997):

“By far the largest known extant population of *B. cartilagineum* occurs in State forest at Little Beach Creek. The rare treeferns, *Cyathea cunninghamii* and *C. marcescens*, also occur at the site. The site is very steep and unsuitable for timber harvesting, and should be reserved, either alone or jointly with the nearby existing Lower Marsh Creek Forest Reserve. There is still the doubt surrounding the *B. cartilagineum* population at the upper Little Beach site as to whether it is sexually reproducing. This trait should be monitored.”

The slender treefern (*Cyathea cunninghamii*) occurs in small numbers at widespread location in Tasmania. The only East Coast populations are in the lower catchments of Lower Marsh and Little Beach creeks in the Little Beach SR and Lower Marsh Creek FR and contain 64 mature plants out of a total estimated population of 250 plants (TSS 2011a). The skirted treefern (*C. Xmarcescens*) is a naturally occurring hybrid between the rare *C. cunninghamii* and the common *C. australis*. This fern is extremely rare with less than 40 plants known from 4 sites, including 34 from the Little Beach SR and Lower Marsh Creek FR where it grows with the *C. cunninghamii* populations (TSS 2011b). These site are the most important location for these very rare and vulnerable treeferns.

The proposed National park also contains some of the most important populations in Tasmania of the rare shrub *Hovea corrickiae*, and the vulnerable fern *Asplenium trichomanes* subsp. *trichomanes*.

Non-vascular flora

Although few surveys for non-vascular flora (mosses, liverworts, lichens) have been undertaken in the area, the diversity of habitats from near-coastal, to dry forests, wet gullies and high elevation rock outcrops would be expected to support a considerable diversity of these plants.

A survey of lichens at South Sister in 2004 identified 178 species, which is a “remarkable level of diversity” and includes one species previously unknown from Tasmania and one species new to science (Kantvilas 2005). As this is only a preliminary survey of small part of the proposed NP, it suggests that the diversity of lichens in the proposed NP would be considerably higher and possibly of regional or statewide significance.

Threatened Fauna

Nine threatened fauna species are reliably known from the proposed National Park and a number of others are potentially present (Table 4, map p. 26).

Table 4. Threatened fauna recorded from proposed National Park and surrounding area.

Name	Common Name	Tas status	National status	Distribution in Proposed Reserve
<i>Accipiter novaehollandiae</i>	grey goshawk	e		Recorded from within the proposed NP, which contains significant areas of wet forest habitat suitable for nesting and foraging.
<i>Aquila audax</i>	wedge-tailed eagle	e	EN	Recorded from within the proposed NP, which contains significant areas of habitat suitable for nesting and foraging. Several known nest sites.
<i>Botaurus poiciloptilus</i>	australasian bittern		EN	Recorded from the region but no accurate records from the proposed NP. Suitable habitat (e.g. floodplain or coastal wetlands) is outside the proposed NP.
<i>Dasyurus maculatus</i> subsp. <i>maculatus</i>	spotted-tailed quoll	r	VU	Recorded from within the proposed NP, which contains significant areas of wet forest habitat.
<i>Haliaeetus leucogaster</i>	white-bellied sea-eagle	v		Recorded from within the proposed NP which contains potential nesting and foraging habitat, wide ranging species more likely to occur on the coast. Nests in Little Beach SR.
<i>Hydroptila scamandra</i>	caddis fly (upper scamander river)	r		Known from the South Esk and Scamander rivers. Unlikely to occur in the proposed NP. Proposed NP protects part of the upper catchment of known habitat.

<i>Lathamus discolor</i>	swift parrot	e	EN	Recorded from within the proposed NP, which contains significant areas of habitat suitable for nesting and foraging (forest with blue gum).
<i>Litoria raniformis</i>	green and golden frog	v	VU	Not recorded from proposed NP where there is limited suitable habitat.
<i>Perameles gunnii</i>	eastern barred bandicoot		VU	Not recorded from proposed NP but likely to occur in suitable habitat (grassy and sedgey forest).
<i>Prototroctes maraena</i>	australian grayling	v	VU	Not recorded from proposed NP and no suitable habitat (occurs in larger rivers).
<i>Sarcophilus harrisii</i>	tasmanian devil	e	EN	Recorded from within the proposed NP, which contains significant areas of habitat suitable for denning and foraging.
<i>Tasmanipatus anophthalmus</i>	blind velvet worm	e	EN	Numerous records from proposed NP which covers a large part of the known habitat of this geographically restricted species.
<i>Tasmanipatus barretti</i>	giant velvet worm	r		Numerous records from proposed NP which covers the southern range limit of this species. Significant areas of suitable habitat.
<i>Thylacinus cynocephalus</i>	thylacine	x	EX	Extinct. No recent records.
<i>Tyto novaehollandiae</i>	masked owl	e	VU	Recorded from within the proposed NP, which contains significant areas of forest habitat suitable for nesting and foraging.

Threatened Species Protection Act 1995 (Tasmania): r = rare, v = vulnerable, e = endangered, x = extinct

Environmental Protection and Biodiversity Conservation Act 1999 (Commonwealth): VU = vulnerable, EN = endangered, CR = critically endangered

Little Creek State Reserve and adjoining land contains the largest stand of blue gum-dominated forest in the northeast and is therefore a key habitat for the endangered swift parrot on the East Coast (map p. 29). Swift parrots nest in colonies in patches of mature forest (at least 100 years old) with abundant tree hollows in proximity to flowering blue gums (*Eucalyptus globulus*) or black gums (*E. ovata*). The proposed National Park and neighbouring coastal strip is perhaps the best example of breeding and foraging habitat for this species in the northern half of the East Coast.

The blind velvet worm (*Tasmanipatus anophthalmus*) is one of the few invertebrates listed as endangered in Tasmania and is restricted to an area of approximately 152 km² around St Marys. The proposed National Park covers most of the core population for this species, which covers an area of only 40 km² around Mount Elephant and the catchments on its eastern slopes (TSS 2005).

According to Dr Peter McQuillan, the blind velvet worm:

“is a native Tasmanian animal of the highest conservation value. Velvet worms in general are of extraordinary scientific interest because they are a conservative, ancient group which shed light on the early evolution of arthropods (insects and

their relatives) and segmented worms. Tasmania is a globally important centre of diversity for this group of animals. The Blind Velvet Worm has one of the most restricted distributions of any species in the world.” (McQuillan 2001)

The giant velvet worm (*T. barretti*) has a larger but still restricted distribution which does not overlap with the blind velvet worm (TSS 2005). The proposed National Park includes significant areas of giant velvet worm habitat at the southern limit of its range. Both species are largely reliant on large old decomposing logs for habitat, particularly in damp places in dry forest such as shady slopes and gullies, but also in wet forest (TSS 2005). Mature forest is ideal habitat because of the abundance of large logs and suitable microclimates.

The proposed National Park would also protect good quality habitat for the wedge-tailed eagle (*Aquila audax fleayi*), masked owl (*Tyto novaehollandiae*) and Tasmanian devil (*Sarcophilus harrisi*).

Vegetation Communities

TASVEG mapping indicates 35 native vegetation communities occur in the proposed National Park, plus small areas of bare rock, regenerating cleared land and easements (Table 5, map p. 23).

Table 5. Mapped vegetation types in the proposed National Park (source: TASVEG 2.0)

TASVEG mapping unit	Hectares
<i>Acacia dealbata</i> forest	538
<i>Acacia melanoxylon</i> forest on rises	2
<i>Acacia melanoxylon</i> swamp forest	1
Agricultural land	24
Broadleaf scrub	54
Dry scrub	2
<i>Eucalyptus amygdalina</i> / <i>E. obliqua</i> damp sclerophyll forest	32
<i>Eucalyptus amygdalina</i> coastal forest and woodland	604
<i>Eucalyptus amygdalina</i> forest and woodland on dolerite	72
<i>Eucalyptus amygdalina</i> forest and woodland on mudstone	677
<i>Eucalyptus brookeriana</i> wet forest	10
<i>Eucalyptus delegatensis</i> dry forest and woodland	1298
<i>Eucalyptus delegatensis</i> forest over rainforest	1
<i>Eucalyptus delegatensis</i> forest with broad-leaf shrubs	472
<i>Eucalyptus delegatensis</i> wet forest (undifferentiated)	198
<i>Eucalyptus globulus</i> dry forest and woodland	84
<i>Eucalyptus globulus</i> wet forest	12
<i>Eucalyptus obliqua</i> dry forest and woodland	269
<i>Eucalyptus obliqua</i> forest over <i>Leptospermum</i>	1
<i>Eucalyptus obliqua</i> forest with broad-leaf shrubs	1416
<i>Eucalyptus obliqua</i> wet forest (undifferentiated)	174
<i>Eucalyptus sieberi</i> forest and woodland not on granite	4856
<i>Eucalyptus sieberi</i> forest and woodland on granite	577
<i>Eucalyptus viminalis</i> - <i>Eucalyptus globulus</i> coastal forest	5
<i>Eucalyptus viminalis</i> grassy forest and woodland	5
Extra-urban miscellaneous	35
Highland grassy sedgeland	9
Highland low rainforest and scrub	1
<i>Leptospermum</i> scrub	6
Lowland grassland complex	3
Lowland sedgy grassland	1
<i>Nothofagus</i> / <i>Atherosperma</i> rainforest	4
Permanent easement	44
Rainforest fernland	84
Regenerating cleared land	5
Rock	27
Silvicultural plantation (unverified)	55
Subalpine heathland	2
Wet heathland	5

Vegetation Communities of Conservation Significance

Significant stands of old growth dry and wet eucalypt forest including communities dominated by *E. amygdalina*, *E. delegatensis*, *E. obliqua* and *E. sieberi*.

Six vegetation types listed as threatened native vegetation communities under the *Nature Conservation Act 2002* are mapped in the proposed Break O'Day National Park, although one community, *Allocasuarina littoralis* forest, has insignificant extent within the proposed reserve where it occurs on private land at the boundary (Table 6, map p. 25).

Table 6. Threatened native vegetation communities in the proposed Break O Day National Park.

Vegetation community	TASVEG code	Hectares
<i>Allocasuarina littoralis</i> forest	NAL	< 1
<i>Eucalyptus brookeriana</i> wet forest	WBR	10.5
<i>Eucalyptus globulus</i> dry forest and woodland	DGL	84
<i>Eucalyptus viminalis</i> - <i>Eucalyptus globulus</i> coastal forest and woodland	DVC	5
Highland grassy sedgeland	MGH	9.5
Rainforest fernland	RFE	84

Brookers gum (*Eucalyptus brookeriana*) wet forest

This forest type is rare in Tasmania and mostly occurs in the far north west. Only around 95 hectares of brookers gum forest occurs in the north east, mostly in small patches scattered across the Ben Lomond bioregion. The 10 hectares of brookers gum forest at Cheeseberry Hill in the proposed National Park is one of the largest patches of this community in the north east and is currently not reserved.

Blue gum (*Eucalyptus globulus*) dry forest and woodland

Dry eucalypt forest dominated by blue gum with a grassy understorey mostly occurs in southeast Tasmania with only limited patches in the northern part of the East Coast as in the proposed National Park and neighbouring near-coastal hills.

White gum (*Eucalyptus viminalis*) and blue gum (*E. globulus*) coastal forest and woodland

This dry coastal forest has a very limited extent in the northeast, mostly confined to a few small patches on private land to the east of the proposed National Park, and extending slightly into the St Patricks Head State Reserve.

Highland grassy sedgeland

This community is dominated by sedges with tussock grasses also present and few trees and shrubs. It occurs on poorly drained flats on the Nicholas Range. Highland grassy sedgeland is most extensive on the Central Plateau and is uncommon in the north east.

Rainforest fernland

This treeless community is dominated by ferns and usually occurs as small patches within rainforest associated with either disturbance events that have eliminated trees or shallow soils. It occurs on steep slopes and in the bottom of gullies on the east side of Mount Elephant. The proposed National Park includes the only occurrences of this community on the East Coast.

Geoconservation

The proposed National Park contains a large number of sites listed on the Tasmanian Geoconservation Database, many of them occurring in the Nicholas Range (Table 7, map p. 30). In addition to the publicly listed sites there is an undisclosed geoconservation feature, a karst site, within the proposed National Park.

Table 7. Sites listed on Tasmanian Geoconservation Database v7.0 within the proposed National Park.

Name	General description	Comments	Significance
St Patricks Head Basaltic Soils	Boundaries as per geological map.	Apparently the only reserved example of Triassic basalt. This basalt has not been dated but is considered Triassic on petrological grounds. (See Huntsmans Ck Triassic Basalt site). Very few basalt soils in Tasmania are unaffected by agriculture - may be important for biodiversity.	Regional
Mt Elephant Karst	Karst features occur in a belt of limestone encircling the flanks of Mt Elephant (St. Marys area), best known in the Gray - Elephant Pass area.	A well developed karst system in Permian limestone. It is the best development of a terrestrial karst system in Permian limestone known in Tasmania. However knowledge of the feature is inadequate for detailed management planning, as the detailed nature of the karst system is poorly known. Area corresponds to Kiernan's (1995) NE34 area.	Tasmania
Huntsmans Creek Waterfall	Near Dublin Town.	A structurally-influenced waterfall on Triassic basalt. Its significance is largely related to the unique nature of the Triassic basalt. (see BOD25)	Local region
St Marys Porphyrite and Catos Creek Dyke	East of St Marys, a large area between Mt Elephant and north of St Marys Pass, and eastwards to the coast. Catos Creek area.	Extrusive Devonian volcanic rocks, overlying Mathinna Group metasediments on an intervening basal breccia. This breccia represents probable slump and landslip deposits produced on the land surface by disturbances immediately prior to volcanic extrusions. The St Marys Porphyrite is the extrusive equivalent of the extensive intrusive granites of the Blue Tier Batholith. It is the only extrusive volcanic rock of Devonian age (388+/-1 Ma) known in Tasmania. Dating of the (unfolded) volcanics is important evidence in dating the underlying folded rocks. See also Little Beach Porphyrite (BOD04).	Tasmania
Mt Nicholas and Blackboy Plains High Plateau Marshes	Various locations including Mt Nicholas summit ridge and Blackboy Plains.	Wetland environments with Holocene (at least) sediment accumulations. They are of potential scientific importance for Holocene stratigraphy, palynology and local wetland ecosystems. Only a few examples have been documented so far, but more undoubtedly exist. Mt Nicholas and Blackboy Plains sites have been listed as separate sites, but are linked to this site in the comments section.	Local

Mt Nicholas High Plateau Marshes	High plateau area of Mt Nicholas summit ridge, north west of St Marys.	Wetland environment with Holocene (at least) sediment accumulations. Of potential scientific importance for Holocene stratigraphy, palynology and local wetland ecosystems. Only a few examples have been documented so far, but more undoubtedly exist. This site is part of the Mt Nicholas and Blackboy Plains High Plateau Marshes (BOD24), in north-eastern Tasmania.	Local
Mt Nicholas Dolerite Residual Peak	Entire Nicholas Range, north-west of St Marys.	Mountain forms with minimal faulting influence - the dominant influences are fluvial and periglacial erosion processes. This site is one of several dolerite residual peaks which are collectively listed under NE Tasmania Dolerite Residual Peaks (FOR02).	Regional
North and South Sister Dolerite Periglacial System	Alpine zones of North and South Sister	Periglacial features include cliffs, tors, stacks, blockstreams, mountain top detritus, talus etc. This site is part of a suite of periglacial mountain sites, which collectively are listed as NE Tasmania Dolerite Periglacial Systems (FOR11).	Regional
Mt Nicholas Dolerite Periglacial System	Alpine Zone of Nicholas Range, north-west of St Marys	Periglacial features including cliffs, tors, stacks, mountain top detritus, talus, blockstreams, slab topples, long-distance blockslides. This site is part of a suite of periglacial mountain sites which are collectively listed as NE Tasmania Dolerite Periglacial Systems (FOR11).	Regional
South Sister Periglacial Boulders and Gelifluction Lobes		The site is significant because it shows that in northeast Tasmania there was intense freeze-thaw and gelifluction to c. 650 m altitude during Oxygen isotope stage 5 (or possibly earlier, if significant surface erosion of the dolerite boulder has occurred and the cosmogenic age is an underestimate).	Regional
North-east Tasmania Dolerite Residual Peaks	Ben Nevis, Mt Saddleback, Mt Victoria, Mt Albert, Mt Young, Mt Blackboy, Mt Nicholas and Tower Hills.	Mountain forms with minimal faulting influence - the dominant influences are fluvial and periglacial erosion processes.	Regional
North-east Tasmania Dolerite Periglacial Systems	Alpine zones of Ben Lomond, ... Mt Nicholas, Tower Hills and North Sister.	Cliffs, tors, stacks, blockstreams, mountain top detritus, talus, slab topples, long-distance blockslides (Mt Victoria) etc.	Regional

National Estate Values

The proposed National Park includes three areas listed on the Australian Government's National Heritage List (DSEWPaC 2011). Little Beach Creek – Lower Marsh Creek Area is listed for its unique rare fern flora and aesthetic values. St Marys Pass State Reserve is significant for the diversity of wet and dry forest types present and the occurrence of rare flora species. St Patricks Head State Reserve is notable for the presence of *Eucalyptus brookeriana* forest, rare flora and a rare example of basalt soils with native vegetation (i.e. not cleared).

The proposed National Park includes areas identified in the Tasmanian Regional Forest Agreement as indicative areas exhibiting limits of range of flora (Scamander – Freycinet: northern limit of many east coast species), indicative areas exhibiting major floral disjunctions (Lower Marsh Creek) and indicative areas for flora species richness and community richness (PLUC 1996). For fauna, the St Marys area is an indicative area for endemic fauna (invertebrates including *Tasmanipatus* spp.) and for relictual fauna. St Marys Pass and Elephant Pass are likely glacial refugia, where some of the only small patches of rainforest and wet forest would have persisted in eastern Tasmania during the last glacial (PLUC 1996).

Naturalness and Restoration

Most of the proposed National Park has a high degree of naturalness. Some areas of State forest have been subject to logging, particularly on the lower slopes of the Nicholas Range and in the Little Beach Creek and Little Marsh Creek catchments east of Mount Elephant. This latter area of State forest has been designated a 'restoration zone' because of the extensive logging that has occurred there. This zone is important for reserve design since it improves the viability and connectivity of the proposed National Park. The regrowth native forest in this zone will gradually increase in structural diversity and condition with minimal management needed. Most of the logged forest is *Eucalyptus sieberi* dry forest which has good potential for recovery. The restoration zone comprises 1035 hectares out of the total 11 664 hectare proposed National Park.

Weeds are a major threat to the integrity of the proposed National Park, particularly near St Marys where a number of serious environmental weeds are well established and are invading native vegetation.

Planning for Climate Change

Even under conservative models of climate change it will have a significant impact on biodiversity. The effects on individual species and communities is difficult to predict.

A CSIRO report offers these principles in regards to the challenge of conserving biodiversity and planning protected areas in a rapidly changing climate (Dunlop & Brown 2008):

Protecting habitat is probably the best way to conserve species under climate change. While the species and ecosystems in any one area will change over time, the greater the total area of habitat available, and the more diverse that habitat, the greater the number of ecosystems and species that will be able to survive. ... Connectivity of habitat at various scales can be important for facilitating the movement of different species, which may increase their viability and ability to respond to climate variability and change.



Steep environmental gradients such as the elevation change from near sea level to the highlands around St Marys (e.g. sea level to 750 metres elevation over 5.5 km at Mount Elephant) and a complex and diverse topography provide a lot more potential for species to migrate and adapt to environmental changes than in a more topographically subdued landscape. The deep gullies in the eastern part of the proposed National Park are likely to function as refugia for fire- and drought-sensitive species because they maintain a cooler wetter climate than the surrounding landscape and provide topographic protection from fire. The proposed National Park is biogeographically significant because it includes the range limits of a number of species and contains several rare and endemic species, consequently it will be an important location for species to migrate or survive in response to climate change.



References

- Bryant, S. L. & Jackson, J. (1999). *Tasmania's Threatened Fauna Handbook*. Threatened Species Unit, Nature Conservation Branch, DPIWE.
- DPIPWE (2011). Natural Values Atlas. <www.naturalvaluesatlas.tas.gov.au>. Accessed 13/6/2011.
- DSEWPaC (2011). Australian Heritage Database. <<http://www.environment.gov.au/heritage/ahdb/index.html>> Accessed 26/8/2011.
- Dunlop, M., and Brown, P. R. (2008) *Implications of Climate Change for Australia's National Reserve System - A Preliminary Assessment*. Report to the Department of Climate Change, and the Department of the Environment, Water, Heritage and the Arts.
- Garrett, M. (1996) *The ferns of Tasmania: their ecology and distribution*. Tasmanian Forest Research Council, Hobart.
- Garrett, M. (1997) *Rare or Threatened Forest Ferns*. A Report to the Tasmanian Regional Forest Agreement. Environment and Heritage Technical Committee, Hobart.
- Harris, S. and Kitchener, A. (2005). *From Forest to Fjaeldmark: Descriptions of Tasmania's Vegetation*. Department of Primary Industries, Water and Environment, Printing Authority of Tasmania. Hobart.
- Jones, D., Wapstra, H., Tonelli, P. & Harris, S. (2000) *The Orchids of Tasmania*. Melbourne University Press, Victoria.
- Kantvilas, G. (2005) 'Preliminary lichen report'. Tasmanian Herbarium, Hobart. <<http://www.southsister.org/articles2/kantvilas.htm>> Accessed 26/8/2011.
- McQuillan, P.B. (2001) 'An assessment of some biodiversity conservation values of Coupe FG122A, St Marys, Tasmania.' Unpublished report, University of Tasmania.
- McQuillan, P.B., Watson, J., Fitzgerald, N.B., Leaman, D. & Obendorf, D. (2009) The importance of ecological processes for terrestrial biodiversity conservation in Tasmania – a review. *Pacific Conservation Biology*, **15**, pp. 171-196.
- NETLT (2009) 'The St Marys Protected Landscape'. North East Tasmania Land Trust Inc. <<http://www.netlandtrust.org.au/project.html>> Accessed 26/8/2011.
- PLUC (1996) *Tasmanian Regional Forest Agreement: National Estate report (Background Report Part H)*. Public Land Use Commission, Hobart.
- TSS (2005) 'Listing statement north-east velvet worms *Tasmanipatus* spp.' Threatened Species Section, DPIWE, Hobart.
- TSS (2011a) Listing Statement for *Cyathea cunninghamii* (slender treefern). Threatened Species Section, Department of Primary Industries, Parks, Water & Environment, Tasmania.
- TSS (2011b) Listing Statement for *Cyathea Xmarcescens* (skirted treefern), Threatened Species Section, Department of Primary Industries, Parks, Water & Environment, Tasmania.

Proposed Break O' Day National Park, northeast Tasmania







-  Proposed National Park
-  Restoration zone



Break O Day National Park proposal v1,
August 2011.
Basemap from TASMALP © State of Tasmania

Land tenure of proposed Break O' Day National Park








-  Proposed Break O' Day NP
-  Major Road
-  Major River

-  Reserve on public land
-  Reserve on private land
-  State forest
-  Crown land

Base layers from theLIST © State of Tasmania.

Geology of proposed Break O' Day National Park




-  Proposed Break O' Day NP
-  Major Road
-  Major River

-  Quaternary - Siliceous Sediments
-  Quaternary - Glacial and Periglacial Features
-  Tertiary - Siliceous Sediments
-  Jurassic - Dolerite
-  Triassic-Jurassic - Basalt
-  Triassic - Carbonaceous Sediments
-  Permian-Triassic - Siliceous Sediments
-  Permian - Calcareous Sediments
-  Devonian - Granite
-  Ordovician-Devonian - Siliceous Sediments

Base layers from theLIST
 © State of Tasmania.
 Geology from Mineral Resources Tasmania
 © State of Tasmania.

Vegetation of proposed Break O' Day National Park














0 1 2 3 4 5 km



Base layers from theLIST © State of Tasmania.
Vegetation: TASVEG 2.0 © State of Tasmania.

- Proposed Break O' Day NP
- Major Road
- Major River




Dry Eucalypt Forest and Woodland

-  DAC Eucalyptus amygdalina coastal forest and woodland
-  DAD Eucalyptus amygdalina forest and woodland on dolerite
-  DAM Eucalyptus amygdalina forest on mudstone
-  DDE Eucalyptus delegatensis dry forest and woodland
-  DGL Eucalyptus globulus dry forest and woodland
-  DOB Eucalyptus obliqua dry forest
-  DVG Eucalyptus viminalis grassy forest and woodland
-  DSC Eucalyptus amygdalina - Eucalyptus obliqua damp sclerophyll forest
-  DSG Eucalyptus sieberi forest and woodland on granite
-  DSO Eucalyptus sieberi forest and woodland not on granite
-  DVC Eucalyptus viminalis-Eucalyptus globulus coastal forest and woodland






Wet Eucalypt forest and woodland

-  WDU Eucalyptus delegatensis wet forest (undifferentiated)
-  WDB Eucalyptus delegatensis forest with broad-leaf shrubs
-  WDR Eucalyptus delegatensis over rainforest
-  WBR Eucalyptus brookeriana wet forest
-  WGL Eucalyptus globulus wet forest
-  WOU Eucalyptus obliqua wet forest (undifferentiated)
-  WOB Eucalyptus obliqua forest with broad-leaf shrubs
-  WOL Eucalyptus obliqua forest over Leptospermum






Rainforest and related scrub

-  RMU Nothofagus rainforest undifferentiated
-  RFE Rainforest fernland
-  RSH Highland low rainforest and scrub


Non-Eucalypt Forest and Woodland

-  NAD Acacia dealbata forest
-  NAF Acacia melanoxylon swamp forest
-  NAL Allocasuarina littoralis forest
-  NAR Acacia melanoxylon forest on rises
-  NLA Leptospermum scoparium-Acacia mucronata forest



Scrub, heathland and coastal complexes

-  SHS Subalpine heathland
-  SHW Wet heathland
-  SBR Broad-leaf scrub
-  SLW Leptospermum scrub
-  SDU Dry scrub






Moorland, Sedgeland, Rushland and Peatland

-  MGH Highland grassy sedgeland

Native Grassland

-  GCL Lowland grassland complex
-  GSL Lowland grassy sedgeland


Agricultural, Urban and Exotic Vegetation




-  FAG Agricultural land
-  FRG Regenerating cleared land
-  FPL Plantations for silviculture
-  FPU
-  FPE Permanent easements
-  FUM Extra-urban miscellaneous


Other Natural Environments

-  ORO Lichen lithosere



Threatened vegetation of proposed Break O' Day National Park




-  Proposed Break O' Day NP
-  Major Road
-  Major River




Dry Eucalypt Forest and Woodland

-  DGL *Eucalyptus globulus* dry forest and woodland
-  DVC *Eucalyptus viminalis*-*Eucalyptus globulus* coastal forest and woodland


Wet Eucalypt forest and woodland

-  WBR *Eucalyptus brookeriana* wet forest

Rainforest and related scrub

-  RFE Rainforest fernland

Non-Eucalypt Forest and Woodland


-  NAL *Allocasuarina littoralis* forest

Moorland, Sedgeland, Rushland and Peatland

-  MGH Highland grassy sedgeland


Base layers from theLIST © State of Tasmania.
Vegetation: TASVEG 2.0 © State of Tasmania.

Threatened fauna records, proposed Break O' Day National Park







- | | | |
|---------------------------------------------------------------------------------------------------------------------|--------------------------------------------|----------------------------------|
| Break O' Day NP 5km buffer | SPECIES | <i>Litoria raniformis</i> |
| Proposed Break O' Day NP | <i>Accipiter novaehollandiae</i> | <i>Perameles gunnii</i> |
| Major Road | <i>Aquila audax subsp. fleayi</i> | <i>Prototroctes maraena</i> |
| Major River | <i>Botaurus poiciloptilus</i> | <i>Sarcophilus harrisii</i> |
| Base layers from theLIST | <i>Dasyurus maculatus subsp. maculatus</i> | <i>Tasmanipatus anophthalmus</i> |
| © State of Tasmania. | <i>Haliaeetus leucogaster</i> | <i>Tasmanipatus barretti</i> |
| Species records from Natural Values Atlas (Tas Govt), only showing records with spatial accuracy under 1100 metres. | <i>Hydroptila scamandra</i> | <i>Tyto novaehollandiae</i> |
| Marine and coastal species not shown. | <i>Lathamus discolor</i> | |

Threatened flora records, proposed Break O' Day National Park





Base layers from theLIST
 © State of Tasmania.
 Species records from Natural Values Atlas
 (Tas Govt), as at July 2011, only showing
 records with spatial accuracy under 1000 m.

-  Proposed Break O' Day NP
-  Break O' Day NP 5km buffer
-  Major Road
-  Major River

SPECIES

- *Acacia siculiformis*
- *Acacia ulicifolia*
- *Asperula subsimplex*
- ▲ *Asplenium trichomanes* subsp. *trichomanes*
- ☆ *Austrostipa nodosa*
- ▲ *Blechnum cartilagineum*
- ◆ *Caladenia caudata*
- ◆ *Caladenia congesta*
- ◆ *Caladenia filamentosa*
- *Calystegia soldanella*
- *Conospermum hookeri*
- ▲ *Cyathea Xmarcescens*
- ▲ *Cyathea cunninghamii*
- *Cynoglossum australe*
- *Desmodium gunnii*
- ☆ *Deyeuxia densa*
- *Euphrasia collina* subsp. *deflexifolia*
- *Glycine microphylla*
- *Hibbertia calycina*
- *Hibbertia virgata*
- ☆ *Hierochloe rariflora*
- *Hovea corrickiae*
- *Hovea longifolia*
- ☆ *Juncus amabilis*
- ☆ *Lachnagrostis billardierei* subsp. *tenuiseta*
- *Lepidium hyssopifolium*
- *Lepidium pseudotasmanicum*
- ★ *Lepidosperma tortuosum*
- *Lotus australis*
- *Melaleuca pustulata*
- *Olearia hookeri*
- ▲ *Pellaea calidirupium*
- *Phebalium daviesii*
- *Pimelea curviflora*
- *Plantago debilis*
- *Pomaderris intermedia*
- *Prostanthera rotundifolia*
- ◆ *Pterostylis atriola*
- ◆ *Pterostylis ziegeleri*
- *Schenkia australis*
- *Scleranthus fasciculatus*
- *Scutellaria humilis*
- ☆ *Sporobolus virginicus*
- *Teucrium corymbosum*
- ◆ *Thelymitra malvina*
- *Thismia rodwayi*
- *Tricoryne elatior*
- *Veronica plebeia*
- *Viola cunninghamii*


Swift parrot habitat, proposed Break O' Day National Park



Base layers from theLIST
 © State of Tasmania.
 Swift parrot habitat mapping from DPIPWE
 (Tas Govt), published 2010, shaded areas
 are parts of swift parrot breeding range that
 have yet to be assessed for habitat value.

- Proposed Break O' Day NP
 - Major Road
 - Major River
- Swift parrot habitat**
- Globulus dominant
 - Globulus sub-dominant
 - Ovata dominant
 - Potential habitat - not mapped

Geoconservation features in proposed Break O' Day National Park



- | | | |
|--------------------------|-------------------------------------------|----------------------------------------------------------|
| Proposed Break O' Day NP | Geoconservation Sites | Mt Nicholas and Blackboy Plains High Plateau Marshes |
| Major Road | Geoconservation sites outside proposed NP | North and South Sister Dolerite Periglacial System |
| Major River | Huntsmans Creek Waterfall | North-east Tasmania Dolerite Periglacial Systems |
| | Mt Elephant Karst | North-east Tasmania Dolerite Residual Peaks |
| | Mt Nicholas Dolerite Periglacial System | South Sister Periglacial Boulders and Gelifluction Lobes |
| | Mt Nicholas Dolerite Residual Peak | St Marys Porphyrite and Catos Creek Dyke |
| | Mt Nicholas High Plateau Marshes | St Patricks Head Basaltic Soils |

Base layers from theLIST © State of Tasmania.
 Geoconservation sites from Tasmanian
 Geoconservation Database v.7.

